

**ADVANCING NEURAL TUBE DEFECT PREVENTION IN
LOW- AND MIDDLE-INCOME COUNTRIES THROUGH
IMPROVED FOLATE STATUS IN WOMEN OF
REPRODUCTIVE AGE —**

**Findings from the Technical *Consultation Folate Status in Women
and Neural Tube Defect Prevention***

REFERENCES

GLOBAL BURDEN OF NEURAL TUBES DEFECTS	1
FOLATE STATUS OF WOMEN OF REPRODUCTIVE AGE AS BASIS OF NTD RISK ASSESSMENT	7
RETHINKING SURVEILLANCE FOR NEURAL TUBE DEFECT PREVENTION: A NEW FRAMEWORK TO ACCELERATE ACTION	13
SAFETY OF FOLIC ACID	15
SHOULD VITAMIN B12 BE CONSIDERED IN ASSESSING NTD RISK.....	20
FRAMEWORK FOR LABORATORY HARMONIZATION OF FOLATE MEASUREMENTS IN LOW-AND-MIDDLE-INCOME COUNTRIES AND REGIONS	29
THE INVESTMENT CASE FOR FOLIC ACID FORTIFICATION.....	34

GLOBAL BURDEN OF NEURAL TUBES DEFECTS

1. Botto LD, Moore CA, Khoury MJ, Erickson JD. Neural-tube defects. *N Engl J Med*. 1999;341(20):1509-19.
2. Bower C, D'Antoine H, Stanley FJ. Neural tube defects in Australia: trends in encephaloceles and other neural tube defects before and after promotion of folic acid supplementation and voluntary food fortification. *Birth Defects Res A Clin Mol Teratol*. 2009;85(4):269-73.
3. WHO/CDC/ICBDSR. Birth defects surveillance: atlas of selected congenital anomalies. Geneva: World Health Organization; 2014. 2014.
4. Parker SE, Yazdy MM, Mitchell AA, Demmer LA, Werler MM. A description of spina bifida cases and co-occurring malformations, 1976-2011. *Am J Med Genet A*. 2014;164A(2):432-40.
5. Christianson AL, Howson CP, Modell B. Global report on birth defects: the hidden toll of dying and disabled children. White Plains (NY): March of Dimes Birth Defects Foundation; 2006. 2006.
6. Zaganjor I, Sekkarie A, Tsang BL, Williams J, Razzaghi H, Mulinare J, et al. Describing the Prevalence of Neural Tube Defects Worldwide: A Systematic Literature Review. *PLoS One*. 2016;11(4):e0151586.
7. Oakley GP, Jr. The scientific basis for eliminating folic acid-preventable spina bifida: a modern miracle from epidemiology. *Ann Epidemiol*. 2009;19(4):226-30.
8. Force USPST, Bibbins-Domingo K, Grossman DC, Curry SJ, Davidson KW, Epling JW, Jr., et al. Folic Acid Supplementation for the Prevention of Neural Tube Defects: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2017;317(2):183-9.
9. Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. MRC Vitamin Study Research Group. *Lancet*. 1991;338(8760):131-7.
10. Smithells RW, Sheppard S, Schorah CJ, Seller MJ, Nevin NC, Harris R, et al. Possible prevention of neural-tube defects by periconceptional vitamin supplementation. *Lancet*. 1980;1(8164):339-40.
11. Czeizel AE, Dudas I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med*. 1992;327(26):1832-5.
12. Viswanathan M, Treiman KA, Kish-Doto J, Middleton JC, Coker-Schwimmer EJ, Nicholson WK. Folic Acid Supplementation for the Prevention of Neural Tube Defects: An Updated Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA*. 2017;317(2):190-203.
13. Agopian AJ, Tinker SC, Lupo PJ, Canfield MA, Mitchell LE, National Birth Defects Prevention S. Proportion of neural tube defects attributable to known risk factors. *Birth Defects Res A Clin Mol Teratol*. 2013;97(1):42-6.
14. Cordero AM, Crider KS, Rogers LM, Cannon MJ, Berry RJ. Optimal serum and red blood cell folate concentrations in women of reproductive age for prevention of neural tube

- defects: World Health Organization guidelines. *MMWR Morb Mortal Wkly Rep.* 2015;64(15):421-3.
15. Crider KS, Devine O, Hao L, Dowling NF, Li S, Molloy AM, et al. Population red blood cell folate concentrations for prevention of neural tube defects: Bayesian model. *BMJ.* 2014;349:g4554.
 16. Williams J, Mai CT, Mulinare J, Isenburg J, Flood TJ, Ethen M, et al. Updated estimates of neural tube defects prevented by mandatory folic Acid fortification - United States, 1995-2011. *MMWR Morb Mortal Wkly Rep.* 2015;64(1):1-5.
 17. De Wals P, Tairou F, Van Allen MI, Uh SH, Lowry RB, Sibbald B, et al. Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med.* 2007;357(2):135-42.
 18. Castilla EE, Orioli IM, Lopez-Camelo JS, Dutra Mda G, Nazer-Herrera J, Latin American Collaborative Study of Congenital M. Preliminary data on changes in neural tube defect prevalence rates after folic acid fortification in South America. *Am J Med Genet A.* 2003;123A(2):123-8.
 19. Castillo-Lancellotti C, Tur JA, Uauy R. Impact of folic acid fortification of flour on neural tube defects: a systematic review. *Public Health Nutr.* 2013;16(5):901-11.
 20. Bhide P, Sagoo GS, Moorthie S, Burton H, Kar A. Systematic review of birth prevalence of neural tube defects in India. *Birth Defects Res A Clin Mol Teratol.* 2013;97(7):437-43.
 21. Moore CA, Li S, Li Z, Hong SX, Gu HQ, Berry RJ, et al. Elevated rates of severe neural tube defects in a high-prevalence area in northern China. *Am J Med Genet.* 1997;73(2):113-8.
 22. Rosenthal J, Casas J, Taren D, Alverson CJ, Flores A, Frias J. Neural tube defects in Latin America and the impact of fortification: a literature review. *Public Health Nutr.* 2014;17(3):537-50.
 23. Rosenthal J, Reeve ME, Ramirez N, Crider KS, Sniezek J, Vellozzi C, et al. Red Blood Cell Folate Insufficiency among nonpregnant Women of Childbearing age in Guatemala 2009 to 2010: Prevalence and predicted Neural Tube Defects risk. *Birth Defects Res A Clin Mol Teratol.* 2016;106(7):587-95.
 24. Susser E, Hoek HW, Brown A. Neurodevelopmental disorders after prenatal famine: The story of the Dutch Famine Study. *Am J Epidemiol.* 1998;147(3):213-6.
 25. Renwick JH. Spina bifida, anencephaly, and potato blight. *Lancet.* 1972;2(7784):967-8.
 26. Smith C, Watt M, Boyd AE, Holmes JC. Anencephaly, spina bifida, and potato blight in the Edinburgh area. *Lancet.* 1973;1(7797):269.
 27. MacMahon B, Yen S, Rothman KJ. Potato blight and neural-tube defects. *Lancet.* 1973;1(7803):598-9.
 28. Lanska DJ. Historical aspects of the major neurological vitamin deficiency disorders: the water-soluble B vitamins. In *Handbook of Clinical Neurology, Vol. 95 (3rd series) - History of Neurology (Chapter 30).* 2010:445-76.

29. Czeizel AE. Nutritional supplementation and prevention of congenital abnormalities. *Curr Opin Obstet Gynecol.* 1995;7(2):88-94.
30. McPartlin J, Halligan A, Scott JM, Darling M, Weir DG. Accelerated folate breakdown in pregnancy. *Lancet.* 1993;341(8838):148-9.
31. Hibbard BM. The Role of Folic Acid in Pregnancy; with Particular Reference to Anaemia, Abruption and Abortion. *J Obstet Gynaecol Br Commonw.* 1964;71:529-42.
32. Elwood JH, Nevin NC. Anencephalus and spina bifida in Belfast (1964-1968). *Ulster Med J.* 1973;42(2):213-22.
33. Frecker M, Fraser FC. Epidemiological studies of neural tube defects in Newfoundland. *Teratology.* 1987;36(3):355-61.
34. Baird PA. Neural tube defects in the Sikhs. *Am J Med Genet.* 1983;16(1):49-56.
35. Verma IC. High frequency of neural-tube defects in North India. *Lancet.* 1978;1(8069):879-80.
36. Kulkarni ML, Mathew MA, Ramachandran B. High incidence of neural-tube defects in South India. *Lancet.* 1987;1(8544):1260.
37. Elwood JH, Scott MJ. Prevalence of anencephalus in the United Kingdom. *Dev Med Child Neurol.* 1982;24(3):394-5.
38. Central Technical Co-ordinating Unit ITC-oUI. Multicentric study of efficacy of periconceptional folic acid containing vitamin supplementation in prevention of open neural tube defects from India. *Indian J Med Res.* 2000;112:206-11.
39. Khoshnood B, Loane M, de Walle H, Arriola L, Addor MC, Barisic I, et al. Long term trends in prevalence of neural tube defects in Europe: population based study. *BMJ.* 2015;351:h5949.
40. Atta CA, Fiest KM, Frolkis AD, Jette N, Pringsheim T, St Germaine-Smith C, et al. Global Birth Prevalence of Spina Bifida by Folic Acid Fortification Status: A Systematic Review and Meta-Analysis. *Am J Public Health.* 2016;106(1):e24-34.
41. Berry RJ, Li Z, Erickson JD, Li S, Moore CA, Wang H, et al. Prevention of neural-tube defects with folic acid in China. China-U.S. Collaborative Project for Neural Tube Defect Prevention. *N Engl J Med.* 1999;341(20):1485-90.
42. Botto LD, Yang Q. 5,10-Methylenetetrahydrofolate reductase gene variants and congenital anomalies: a HuGE review. *Am J Epidemiol.* 2000;151(9):862-77.
43. Northrup H, Volcik KA. Spina bifida and other neural tube defects. *Curr Probl Pediatr.* 2000;30(10):313-32.
44. Cawley S, Mullaney L, McKeating A, Farren M, McCartney D, Turner MJ. Knowledge about folic acid supplementation in women presenting for antenatal care. *Eur J Clin Nutr.* 2016;70(11):1285-90.
45. Cawley S, Mullaney L, McKeating A, Farren M, McCartney D, Turner MJ. A review of European guidelines on periconceptional folic acid supplementation. *Eur J Clin Nutr.* 2016;70(2):143-54.

46. Friberg AK, Jorgensen FS. Few Danish pregnant women follow guidelines on periconceptional use of folic acid. *Dan Med J*. 2015;62(3).
47. Alasfoor D, Elsayed MK, Mohammed AJ. Spina bifida and birth outcome before and after fortification of flour with iron and folic acid in Oman. *East Mediterr Health J*. 2010;16(5):533-8.
48. Afshar M, Golalipour MJ, Farhud D. Epidemiologic aspects of neural tube defects in South East Iran. *Neurosciences (Riyadh)*. 2006;11(4):289-92.
49. Kato K, Fujiki K. Incidence of congenital malformations in Tokyo Metropolitan Hospitals, 1979-1993. *Brain Dev*. 1996;18(3):230-3.
50. McDonnell R, Delany V, O'Mahony MT, Mullaney C, Lee B, Turner MJ. Neural tube defects in the Republic of Ireland in 2009-11. *J Public Health (Oxf)*. 2015;37(1):57-63.
51. Kulkarni ML, Mathew MA, Reddy V. The range of neural tube defects in southern India. *Arch Dis Child*. 1989;64(2):201-4.
52. Airede KI. Neural tube defects in the middle belt of Nigeria. *J Trop Pediatr*. 1992;38(1):27-30.
53. Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths: rates, risk factors, and acceleration towards 2030. *Lancet*. 2016;387(10018):587-603.
54. Caudill MA, Cruz AC, Gregory JF, 3rd, Hutson AD, Bailey LB. Folate status response to controlled folate intake in pregnant women. *J Nutr*. 1997;127(12):2363-70.
55. Daly LE, Kirke PN, Molloy A, Weir DG, Scott JM. Folate levels and neural tube defects. Implications for prevention. *JAMA*. 1995;274(21):1698-702.
56. Laurence KM. Effect of early surgery for spina bifida cystica on survival and quality of life. *Lancet*. 1974;1(7852):301-4.
57. Shurtleff DB, Luthy DA, Nyberg DA, Benedetti TJ, Mack LA. Meningomyelocele: management in utero and post natum. *Ciba Found Symp*. 1994;181:270-80; discussion 80-6.
58. Hunt GM, Oakeshott P. Lifestyle in adults aged 35 years who were born with open spina bifida: prospective cohort study. *Cerebrospinal Fluid Res*. 2004;1(1):4.
59. Wong LY, Paulozzi LJ. Survival of infants with spina bifida: a population study, 1979-94. *Paediatr Perinat Epidemiol*. 2001;15(4):374-8.
60. Shin M, Kucik JE, Siffel C, Lu C, Shaw GM, Canfield MA, et al. Improved survival among children with spina bifida in the United States. *J Pediatr*. 2012;161(6):1132-7.
61. Bowman RM, McLone DG, Grant JA, Tomita T, Ito JA. Spina bifida outcome: a 25-year prospective. *Pediatr Neurosurg*. 2001;34(3):114-20.
62. Sutton LN. Fetal surgery for neural tube defects. *Best Pract Res Clin Obstet Gynaecol*. 2008;22(1):175-88.
63. Oakeshott P, Hunt GM, Poulton A, Reid F. Expectation of life and unexpected death in open spina bifida: a 40-year complete, non-selective, longitudinal cohort study. *Dev Med Child Neurol*. 2010;52(8):749-53.

64. Wang Y, Hu J, Druschel CM, Kirby RS. Twenty-five-year survival of children with birth defects in New York State: a population-based study. *Birth Defects Res A Clin Mol Teratol.* 2011;91(12):995-1003.
65. Kancherla V, Druschel CM, Oakley GP, Jr. Population-based study to determine mortality in spina bifida: New York State Congenital Malformations Registry, 1983 to 2006. *Birth Defects Res A Clin Mol Teratol.* 2014;100(8):563-75.
66. Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE, et al. Global, regional, and national causes of child mortality in 2000-13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet.* 2015;385(9966):430-40.
67. Lamberti LM, Ashraf S, Walker CL, Black RE. A Systematic Review of the Effect of Rotavirus Vaccination on Diarrhea Outcomes Among Children Younger Than 5 Years. *Pediatr Infect Dis J.* 2016;35(9):992-8.
68. Lawn JE, Blencowe H, Pattinson R, Cousens S, Kumar R, Ibiebele I, et al. Stillbirths: Where? When? Why? How to make the data count? *Lancet.* 2011;377(9775):1448-63.
69. Allagh KP, Shamanna BR, Murthy GV, Ness AR, Doyle P, Neogi SB, et al. Birth prevalence of neural tube defects and orofacial clefts in India: a systematic review and meta-analysis. *PLoS One.* 2015;10(3):e0118961.
70. Cousens S, Blencowe H, Stanton C, Chou D, Ahmed S, Steinhardt L, et al. National, regional, and worldwide estimates of stillbirth rates in 2009 with trends since 1995: a systematic analysis. *Lancet.* 2011;377(9774):1319-30.
71. Sutton M, Daly LE, Kirke PN. Survival and disability in a cohort of neural tube defect births in Dublin, Ireland. *Birth Defects Res A Clin Mol Teratol.* 2008;82(10):701-9.
72. Blencowe H, Calvert Ph DC, Lawn JE, Cousens S, Campbell OM. Measuring maternal, foetal and neonatal mortality: Challenges and solutions. *Best Pract Res Clin Obstet Gynaecol.* 2016;36:14-29.
73. Haase J, Green A, Hauge M, Holm NV, Mathiasen H. A cohort study of neural tube defects (NTD) in Denmark covering the first seven years of life. *Childs Nerv Syst.* 1987;3(2):117-20.
74. Kalucy M, Bower C, Stanley F, Burton P. Survival of infants with neural tube defects in Western Australia 1966-1990. *Paediatr Perinat Epidemiol.* 1994;8(3):334-51.
75. McDonnell GV, McCann JP. Why do adults with spina bifida and hydrocephalus die? A clinic-based study. *Eur J Pediatr Surg.* 2000;10 Suppl 1:31-2.
76. Preis K, Swiatkowska-Freund M, Janczewska I. Spina bifida--a follow-up study of neonates born from 1991 to 2001. *J Perinat Med.* 2005;33(4):353-6.
77. Roach JW, Short BF, Saltzman HM. Adult consequences of spina bifida: a cohort study. *Clin Orthop Relat Res.* 2011;469(5):1246-52.
78. Rowland CA, Correa A, Cragan JD, Alverson CJ. Are encephaloceles neural tube defects? *Pediatrics.* 2006;118(3):916-23.

79. Cortes F, Mellado C, Pardo RA, Villarroel LA, Hertrampf E. Wheat flour fortification with folic acid: changes in neural tube defects rates in Chile. *Am J Med Genet A*. 2012;158A(8):1885-90.
80. Sayed AR, Bourne D, Pattinson R, Nixon J, Henderson B. Decline in the prevalence of neural tube defects following folic acid fortification and its cost-benefit in South Africa. *Birth Defects Res A Clin Mol Teratol*. 2008;82(4):211-6.
81. Hertrampf E, Cortes F. Folic acid fortification of wheat flour: Chile. *Nutr Rev*. 2004;62(6 Pt 2):S44-8; discussion S9.
82. Lopez-Camelo JS, Orioli IM, da Graca Dutra M, Nazer-Herrera J, Rivera N, Ojeda ME, et al. Reduction of birth prevalence rates of neural tube defects after folic acid fortification in Chile. *Am J Med Genet A*. 2005;135(2):120-5.
83. Pacheco SS, Braga C, Souza AI, Figueiroa JN. Effects of folic acid fortification on the prevalence of neural tube defects. *Rev Saude Publica*. 2009;43(4):565-71.
84. Ricks DJ, Rees CA, Osborn KA, Crookston BT, Leaver K, Merrill SB, et al. Peru's national folic acid fortification program and its effect on neural tube defects in Lima. *Rev Panam Salud Publica*. 2012;32(6):391-8.
85. Santos LM, Lecca RC, Cortez-Escalante JJ, Sanchez MN, Rodrigues HG. Prevention of neural tube defects by the fortification of flour with folic acid: a population-based retrospective study in Brazil. *Bull World Health Organ*. 2016;94(1):22-9.
86. Lo A, Polsek D, Sidhu S. Estimating the burden of neural tube defects in low- and middle-income countries. *J Glob Health*. 2014;4(1):010402.

FOLATE STATUS OF WOMEN OF REPRODUCTIVE AGE AS BASIS OF NTD RISK ASSESSMENT

1. Daly, L.E., et al. Folate levels and neural tube defects. Implications for prevention. *JAMA*, 1995. 274(21): p. 1698-702.
2. World Health Organization. Nutrition for Health and Development, Guideline. Optimal serum and red blood cell folate concentrations in women of reproductive age for prevention of neural tube defects. 2015, World Health Organization,: Geneva, Switzerland. p. 1 online resource (1 PDF file (viii, 38 pages)).
3. Pfeiffer, C.M., et al. Comparison of serum and red blood cell folate microbiologic assays for national population surveys. *J Nutr*, 2011. 141(7): p. 1402-9.
4. Organization, W.H. Serum and red blood cell folate concentrations for assessing folate status in populations, V.a.M.N.I. System, Editor. 2012, World Health Organization: Geneva.
5. Bailey, L.B., et al. Biomarkers of Nutrition for Development-Folate Review. *J Nutr*, 2015. 145(7): p. 1636S-1680S.
6. Bailey, L. and M. Caudill. Folate, in *Present Knowledge in Nutrition*, 10th Edition, J. Erdman, I. MacDonald, and S. Zeisel, Editors. 2012, Wiley-Blackwell: Hoboken, New Jersey. p. 321-342.
7. Hao, L., et al. Geographical, seasonal and gender differences in folate status among Chinese adults. *J Nutr*, 2003. 133(11): p. 3630-5.
8. Pfeiffer, C.M., et al. Estimation of trends in serum and RBC folate in the U.S. population from pre- to postfortification using assay-adjusted data from the NHANES 1988-2010. *J Nutr*, 2012. 142(5): p. 886-93.
9. Yang, Q., et al. Folic acid source, usual intake, and folate and vitamin B-12 status in US adults: National Health and Nutrition Examination Survey (NHANES) 2003-2006. *Am J Clin Nutr*, 2010. 91(1): p. 64-72.
10. Berry, R.J., et al. Prevention of neural-tube defects with folic acid in China. China-U.S. Collaborative Project for Neural Tube Defect Prevention. *N Engl J Med*, 1999. 341(20): p. 1485-90.
11. Yeung, L., Q. Yang, and R.J. Berry. Contributions of total daily intake of folic acid to serum folate concentrations. *JAMA*, 2008. 300(21): p. 2486-7.
12. Hertrampf, E., et al. Consumption of folic acid-fortified bread improves folate status in women of reproductive age in Chile. *J Nutr*, 2003. 133(10): p. 3166-9.
13. Hopkins, S.M., et al. Impact of voluntary fortification and supplement use on dietary intakes and biomarker status of folate and vitamin B-12 in Irish adults. *Am J Clin Nutr*, 2015. 101(6): p. 1163-72.
14. Caudill, M.A., et al. Folate status response to controlled folate intake in pregnant women. *J Nutr*, 1997. 127(12): p. 2363-70.

15. Hao, L., et al. Folate status and homocysteine response to folic acid doses and withdrawal among young Chinese women in a large-scale randomized double-blind trial. *Am J Clin Nutr*, 2008. 88(2): p. 448-57.
16. Tighe, P., et al. A dose-finding trial of the effect of long-term folic acid intervention: implications for food fortification policy. *Am J Clin Nutr*, 2011. 93(1): p. 11-8.
17. Duffy, M.E., et al. Biomarker responses to folic acid intervention in healthy adults: a meta-analysis of randomized controlled trials. *Am J Clin Nutr*, 2014. 99(1): p. 96-106.
18. Wright, A.J., et al. Comparison of (6 S)-5-methyltetrahydrofolic acid v. folic acid as the reference folate in longer-term human dietary intervention studies assessing the relative bioavailability of natural food folates: comparative changes in folate status following a 16-week placebo-controlled study in healthy adults. *Br J Nutr*, 2010. 103(5): p. 724-9.
19. Pfeiffer, C.M., et al. Applying inappropriate cutoffs leads to misinterpretation of folate status in the US population. *Am J Clin Nutr*, 2016. 104(6): p. 1607-1615.
20. Crider, K.S., et al. Population red blood cell folate concentrations for prevention of neural tube defects: Bayesian model. *BMJ*, 2014. 349: p. g4554.
21. Shane, B. Folate Chemistry and Metabolism, in *Folate in Health and Disease*, L. Bailey, Editor. 2010, CRC Press, Taylor Francis Group: Boca Raton, FL. p. 1-24.
22. Mason, J.B. Biomarkers of nutrient exposure and status in one-carbon (methyl) metabolism. *J Nutr*, 2003. 133 Suppl 3: p. 941S-947S.
23. Clifford, A.J., et al. Erythrocyte folate and its response to folic acid supplementation is assay dependent in women. *J Nutr*, 2005. 135(1): p. 137-43.
24. Houghton, L.A., et al. Long-term effect of low-dose folic acid intake: potential effect of mandatory fortification on the prevention of neural tube defects. *Am J Clin Nutr*, 2011. 94(1): p. 136-41.
25. Wu, A., et al. Folate deficiency in the alcoholic--its relationship to clinical and haematological abnormalities, liver disease and folate stores. *Br J Haematol*, 1975. 29(3): p. 469-78.
26. Drogan, D., et al. Plasma folate as marker of folate status in epidemiological studies: the European Investigation into Cancer and Nutrition (EPIC)-Potsdam study. *Br J Nutr*, 2004. 92(3): p. 489-96.
27. Rosenthal, J., et al. Folate deficiency prevalence in women of childbearing age in Belize and negatively impacted by coexisting vitamin B12 deficiency: Belize National Micronutrient Survey 2011. *The Journal of Nutrition*, In press.
28. Molloy, A.M., et al. Maternal vitamin B12 status and risk of neural tube defects in a population with high neural tube defect prevalence and no folic Acid fortification. *Pediatrics*, 2009. 123(3): p. 917-23.
29. Ueland, P.M., et al. Biological and clinical implications of the MTHFR C677T polymorphism. *Trends Pharmacol Sci*, 2001. 22(4): p. 195-201.

30. Crider, K.S., et al. MTHFR 677C->T genotype is associated with folate and homocysteine concentrations in a large, population-based, double-blind trial of folic acid supplementation. *Am J Clin Nutr*, 2011. 93(6): p. 1365-72.
31. Clarke, R., et al. Homocysteine and coronary heart disease: meta-analysis of MTHFR case-control studies, avoiding publication bias. *PLoS Med*, 2012. 9(2): p. e1001177.
32. Yang, Q.H., et al. Prevalence and effects of gene-gene and gene-nutrient interactions on serum folate and serum total homocysteine concentrations in the United States: findings from the third National Health and Nutrition Examination Survey DNA Bank. *American Journal of Clinical Nutrition*, 2008. 88(1): p. 232-246.
33. Holmes, M.V., et al. Effect modification by population dietary folate on the association between MTHFR genotype, homocysteine, and stroke risk: a meta-analysis of genetic studies and randomised trials. *Lancet*, 2011. 378(9791): p. 584-94.
34. Yan, L., et al. Association of the maternal MTHFR C677T polymorphism with susceptibility to neural tube defects in offsprings: evidence from 25 case-control studies. *PLoS One*, 2012. 7(10): p. e41689.
35. Zhang, T., et al. Genetic variants in the folate pathway and the risk of neural tube defects: a meta-analysis of the published literature. *PLoS One*, 2013. 8(4): p. e59570.
36. Williams, J., et al. Updated estimates of neural tube defects prevented by mandatory folic Acid fortification - United States, 1995-2011. *MMWR Morb Mortal Wkly Rep*, 2015. 64(1): p. 1-5.
37. Ganji, V. and M.R. Kafai. Trends in serum folate, RBC folate, and circulating total homocysteine concentrations in the United States: analysis of data from National Health and Nutrition Examination Surveys, 1988-1994, 1999-2000, and 2001-2002. *J Nutr*, 2006. 136(1): p. 153-8.
38. Centers for Disease Control and Prevention (U.S.), National Health and Nutrition Examination Survey (U.S.), and National Center for Environmental Health (U.S.). Division of Laboratory Sciences. Second National Report on Biochemical Indicators of Diet and Nutrition in the U.S. Population. 2012, Atlanta, Ga.: National Center for Environmental Health, The Centers for Disease Control and Prevention, Division of Laboratory Sciences. 484 pages.
39. Botto, L.D. and Q. Yang. 5,10-Methylenetetrahydrofolate reductase gene variants and congenital anomalies: a HuGE review. *Am J Epidemiol*, 2000. 151(9): p. 862-77.
40. Giles, W.H., et al. Thermolabile methylenetetrahydrofolate reductase polymorphism (C677T) and total homocysteine concentration among African-American and white women. *Ethn Dis*, 1998. 8(2): p. 149-57.
41. Metz, J. Folic acid metabolism and malaria. *Food Nutr Bull*, 2007. 28(4 Suppl): p. S540-9.
42. Nzila, A., J. Okombo, and A.M. Molloy. Impact of folate supplementation on the efficacy of sulfadoxine/pyrimethamine in preventing malaria in pregnancy: the potential of 5-methyl-tetrahydrofolate. *J Antimicrob Chemother*, 2014. 69(2): p. 323-30.
43. Gregson, A. and C.V. Plowe. Mechanisms of resistance of malaria parasites to antifolates. *Pharmacol Rev*, 2005. 57(1): p. 117-45.

44. Oppenheimer, S.J. and P. Cashin. Serum and red cell folate levels associated with malarial parasitaemia. *Trans R Soc Trop Med Hyg*, 1986. 80(1): p. 169-71.
45. Chango, A. and L. Abdennebi-Najar. Folate metabolism pathway and *Plasmodium falciparum* malaria infection in pregnancy. *Nutr Rev*, 2011. 69(1): p. 34-40.
46. Mazumdar, M., et al. Arsenic is associated with reduced effect of folic acid in myelomeningocele prevention: a case control study in Bangladesh. *Environ Health*, 2015. 14: p. 34.
47. Kamynina, E., et al. Arsenic trioxide targets MTHFD1 and SUMO-dependent nuclear de novo thymidylate biosynthesis. *Proc Natl Acad Sci U S A*, 2017. 114(12): p. E2319-E2326.
48. Howe, C.G., et al. Folate and cobalamin modify associations between S-adenosylmethionine and methylated arsenic metabolites in arsenic-exposed Bangladeshi adults. *J Nutr*, 2014. 144(5): p. 690-7.
49. Hall, M.N. and M.V. Gamble. Nutritional manipulation of one-carbon metabolism: effects on arsenic methylation and toxicity. *J Toxicol*, 2012. 2012: p. 595307.
50. Suarez, L., et al. Neural tube defects on the Texas-Mexico border: what we've learned in the 20 years since the Brownsville cluster. *Birth Defects Res A Clin Mol Teratol*, 2012. 94(11): p. 882-92.
51. Voss, K.A., R.T. Riley, and J. Gelineau-van Waes. Fumonisin B₁ induced neural tube defects were not increased in LM/Bc mice fed folate-deficient diet. *Mol Nutr Food Res*, 2014. 58(6): p. 1190-8.
52. Gao, L.J., et al. Maternal overweight and obesity and the risk of neural tube defects: a case-control study in China. *Birth Defects Res A Clin Mol Teratol*, 2013. 97(3): p. 161-5.
53. Ray, J.G., et al. Greater maternal weight and the ongoing risk of neural tube defects after folic acid flour fortification. *Obstet Gynecol*, 2005. 105(2): p. 261-5.
54. Mojtabai, R. Body mass index and serum folate in childbearing age women. *Eur J Epidemiol*, 2004. 19(11): p. 1029-36.
55. Tinker, S.C., et al. Does obesity modify the association of supplemental folic acid with folate status among nonpregnant women of childbearing age in the United States? *Birth Defects Res A Clin Mol Teratol*, 2012. 94(10): p. 749-55.
56. Rasmussen, S.A., et al. Maternal obesity and risk of neural tube defects: a metaanalysis. *Am J Obstet Gynecol*, 2008. 198(6): p. 611-9.
57. Pfeiffer, C.M., et al. Dietary Supplement Use and Smoking Are Important Correlates of Biomarkers of Water-Soluble Vitamin Status after Adjusting for Sociodemographic and Lifestyle Variables in a Representative Sample of U.S. Adults. *J Nutr*, 2013. 143(6): p. 957S-65S.
58. Rosenthal, J., et al. Red Blood Cell Folate Insufficiency among nonpregnant Women of Childbearing age in Guatemala 2009 to 2010: Prevalence and predicted Neural Tube Defects risk. *Birth Defects Res A Clin Mol Teratol*, 2016. 106(7): p. 587-95.

59. Shahab-Ferdows, S., et al. Regional, Socioeconomic, and Dietary Risk Factors for Vitamin B-12 Deficiency Differ from Those for Folate Deficiency in Cameroonian Women and Children. *J Nutr*, 2015. 145(11): p. 2587-95.
60. Engle-Stone, R., et al. Stunting prevalence, plasma zinc concentrations, and dietary zinc intakes in a nationally representative sample suggest a high risk of zinc deficiency among women and young children in Cameroon. *J Nutr*, 2014. 144(3): p. 382-91.
61. Imhoff-Kunsch, B., et al., Wheat flour fortification is unlikely to benefit the neediest in Guatemala. *J Nutr*, 2007. 137(4): p. 1017-22.
62. Popkin, B.M. and P.S. Haines. Factors affecting food selection: the role of economics. *J Am Diet Assoc*, 1981. 79(4): p. 419-25.
63. Iannotti, L.L., et al. Food prices and poverty negatively affect micronutrient intakes in Guatemala. *J Nutr*, 2012. 142(8): p. 1568-76.
64. Frohlich, K.L. and L. Potvin. Transcending the known in public health practice: the inequality paradox: the population approach and vulnerable populations. *Am J Public Health*, 2008. 98(2): p. 216-21.
65. Khan, S., et al. Health implications of food patterns in Belize. *Topics in Clinical Nutrition*, 2009. 24: p. 73-81.
66. Cosminsky, S. Women and Health Care on a Guatemalan plantation. *Social Science and Medicine*, 1987. 25: p. 1163-1173.
67. Tsang, B.L., et al. Assessing the association between the methylenetetrahydrofolate reductase (MTHFR) 677C>T polymorphism and blood folate concentrations: a systematic review and meta-analysis of trials and observational studies. *Am J Clin Nutr*, 2015. 101(6): p. 1286-94.
68. Marchetta, C.M., et al. Assessing the association between natural food folate intake and blood folate concentrations: a systematic review and Bayesian meta-analysis of trials and observational studies. *Nutrients*, 2015. 7(4): p. 2663-86.
69. Branum, A.M., R. Bailey, and B.J. Singer, Dietary supplement use and folate status during pregnancy in the United States. *J Nutr*, 2013. 143(4): p. 486-92.
70. Parker, S.E., et al. Updated National Birth Prevalence estimates for selected birth defects in the United States, 2004-2006. *Birth Defects Res A Clin Mol Teratol*, 2010. 88(12): p. 1008-16.
71. Centers for Disease Control and Prevention (CDC), Grand Rounds: additional opportunities to prevent neural tube defects with folic acid fortification. *MMWR Morb Mortal Wkly Rep*, 2010. 59(31): p. 980-4
72. Group, M.V.S.R., Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. *Lancet*, 1991. 338(8760): p. 131-7.
73. Czeizel, A.E. and I. Dudás. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med*, 1992. 327(26): p. 1832-5.
74. Cortés, F., et al. Wheat flour fortification with folic acid: changes in neural tube defects rates in Chile. *Am J Med Genet A*, 2012. 158A(8): p. 1885-90.

75. De Wals, P., et al. Reduction in neural-tube defects after folic acid fortification in Canada. *N Engl J Med*, 2007. 357(2): p. 135-42.
76. De Wals, P., et al., Spina bifida before and after folic acid fortification in Canada. *Birth Defects Res A Clin Mol Teratol*, 2008. 82(9): p. 622-6.
77. Chen, L.T. and M.A. Rivera. The Costa Rican experience: reduction of neural tube defects following food fortification programs. *Nutr Rev*, 2004. 62(6 Pt 2): p. S40-3.
78. Williams, L.J., et al. Prevalence of spina bifida and anencephaly during the transition to mandatory folic acid fortification in the United States. *Teratology*, 2002. 66(1): p. 33-9.
79. Sayed, A.R., et al. Decline in the prevalence of neural tube defects following folic acid fortification and its cost-benefit in South Africa. *Birth Defects Res A Clin Mol Teratol*, 2008. 82(4): p. 211-6.
80. Tinker, S.C., et al. U.S. women of childbearing age who are at possible increased risk of a neural tube defect-affected pregnancy due to suboptimal red blood cell folate concentrations, National Health and Nutrition Examination Survey 2007 to 2012. *Birth Defects Res A Clin Mol Teratol*, 2015. 103(6): p. 517-26.
81. CDC Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. *Morbidity and Mortality Weekly Report*, 1992.
82. Molloy, A.M. and J.M. Scott. Microbiologic assay for serum, plasma, and red cell folate using cryopreserved, microtiter plate method. *Methods Enzymol*, 1997. 281: p. 43-53.
83. Serdula, M.K., et al. Micronutrient status in Jordan: 2002 and 2010. *Eur J Clin Nutr*, 2014. 68(10): p. 1124-8.
84. Labadarios, D., et al. Executive summary of the National Food Consumption Survey Fortification Baseline (NFCS-FB-I). *South African Journal of Clinical Nutrition*, 2008. 21(3 (Suppl 2)): p. 247-300.
85. Freire, W., et al. Tomo I: Encuesta Nacional de Salud y Nutricion de la poblacion ecuatoriana de cero a 59 anos. ENSANUT-ECU 2012, M.d.S.P.I.N.d.E.y. Censos, Editor. 2014: Quito-Ecuador.
86. Rosenthal, J., et al. Folate and Vitamin B12 Deficiency Among Non-pregnant Women of Childbearing-Age in Guatemala 2009-2010: Prevalence and Identification of Vulnerable Populations. *Matern Child Health J*, 2015. 19(10): p. 2272-85.
87. McLean, E., B. de Benoist, and L.H. Allen. Review of the magnitude of folate and vitamin B12 deficiencies worldwide. *Food Nutr Bull*, 2008. 29(2 Suppl): p. S38-51.
88. Institute of Medicine. Dietary Reference Intakes for thiamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin and choline. 1998, National Academy Press: Washington (DC).

RETHINKING SURVEILLANCE FOR NEURAL TUBE DEFECT PREVENTION: A NEW FRAMEWORK TO ACCELERATE ACTION

1. Christianson AL, Howson CP, Modell B. Global report on birth defects: the hidden toll of dying and disabled children. White Plains (NY): March of Dimes Defects Foundation; 2006.
2. Zaganjor I, Sekkarie A, Tsang BL, Williams J, Razzaghi H, Mulinare J, et al. Describing the Prevalence of Neural Tube Defects Worldwide: A Systematic Literature Review. *PLoS One*. 2016;11(4):e0151586.
3. Botto LD, Moore CA, Khoury MJ, Erickson JD. Neural-tube defects. *N Engl J Med*. 1999;341(20):1509-1519.
4. Khoshnood B, Loane M, de Walle H, Arriola L, Addor MC, Barisic I, et al. Long term trends in prevalence of neural tube defects in Europe: population based study. *Br Med J*. 2015;351:h5949.
5. GAIN, Micronutrient Forum, Sight&Life. The #Futurefortified global summit on food fortification - Events proceedings and recommendations for food fortification programs. July, 2016.
6. Thacker SB, Berkelman R. *MMWR*. 2012;Suppl 61(03):10-14.
7. McDonnell R, Delany V, O'Mahony MT, Mullaney C, Lee B, Turner MJ. Neural tube defects in the Republic of Ireland in 2009-11. *J Public Health (Oxf)*. 2015;37(1):57-63.
8. Bowman RM, McLone DG, Grant JA, Tomita T, Ito JA. Spina bifida outcome: a 25-year prospective. *Pediatr Neurosurg*. 2001;34(3):114-120.
9. Haase J, Green A, Hauge M, Holm NV, Mathiasen H. A cohort study of neural tube defects (NTD) in Denmark covering the first seven years of life. *Childs Nerv Syst*. 1987;3(2):117-120.
10. Hunt GM, Oakeshott P. Lifestyle in adults aged 35 years who were born with open spina bifida: prospective cohort study. *Cerebrospinal Fluid Res*. 2004;1(1):4.
11. Kalucy M, Bower C, Stanley F, Burton P. Survival of infants with neural tube defects in Western Australia 1966-1990. *Paediatr Perinat Epidemiol*. 1994;8(3):334-351.
12. Kancherla V, Druschel CM, Oakley GP Jr. Population-based study to determine mortality in spina bifida: New York State Congenital Malformations Registry, 1983 to 2006. *Birth Defects Res A Clin Mol Teratol*. 2014;100(8):563-575.
13. McDonnell GV, McCann JP. Why do adults with spina bifida and hydrocephalus die? A clinic-based study. *Eur J Pediatr Surg*. 2000;10(Supple 1):31-32.
14. Preis K, Swiatkowska-Freund M, Janczewska I. Spina bifida--a follow-up study of neonates born from 1991 to 2001. *J Perinat Med*. 2005;33(4):353-356.

15. Roach JW, Short BF, Saltzman HM. Adult consequences of spina bifida: a cohort study. *Clin Orthop Relat Res.* 2011;469(5):1246-1252.
16. Shin M, Kucik JE, Siffel C, Lu C, Shaw GM, Canfield MA, et al. Improved Survival Among Children with Spina Bifida in the United States. *J Pediatr.* 2012;161(6):1132-1137.
17. Wang Y, Hu J, Druschel CM, Kirby RS. Twenty-five-year survival of children with birth defects in New York State: a population-based study. *Birth Defects Res A Clin Mol Teratol.* 2011;91(12):995-1003.
18. Wong LY, Paulozzi LJ. Survival of infants with spina bifida: a population study, 1979-94. *Paediatr Perinat Epidemiol.* 2001;15(4):374-378.
19. Cordero AM, Crider KS, Rogers LM, Cannon MJ, Berry RJ. Optimal serum and red blood cell folate concentrations in women of reproductive age for prevention of neural tube defects: World Health Organization guidelines. *MMWR Morb Mortal Wkly Rep.* 2015;64(15):421-423.
20. Crider KS, Devine O, Hao L, Dowling NF, Li S, Molloy AM, et al. Population red blood cell folate concentrations for prevention of neural tube defects: Bayesian model. *Br Med J.* 2014;349:g4554.
21. Daly LE, Kirke PN, Molloy A, Weir DG, Scott JM. Folate levels and neural tube defects. Implications for prevention. *JAMA.* 1995;274(21):1698-1702.
22. Rosenthal J, Reeve ME, Ramirez N, Crider KS, Sniezek J, Vellozzi C, et al. Red Blood Cell Folate Insufficiency among nonpregnant Women of Childbearing age in Guatemala 2009 to 2010: Prevalence and predicted Neural Tube Defects risk. *Birth Defects Res A Clin Mol Teratol.* 2016;106(7):587-595.
23. Farrel, et al. 2003.
24. Calvo EB, Biblieri A. Impact of folic acid fortification on women's nutritional status and on the prevalence of neural tube defects. *Arch Arg Ped.* 2008;106(6):492-298.

SAFETY OF FOLIC ACID

1. Force USPST, et al. (2017) Folic Acid Supplementation for the Prevention of Neural Tube Defects: US Preventive Services Task Force Recommendation Statement. *JAMA* 317(2):183-189.
2. Gomes S, Lopes C, & Pinto E (2016) Folate and folic acid in the periconceptional period: recommendations from official health organizations in thirty-six countries worldwide and WHO. *Public Health Nutr* 19(1):176-189.
3. Moussa HN, Hosseini Nasab S, Haidar ZA, Blackwell SC, & Sibai BM (2016) Folic acid supplementation: what is new? Fetal, obstetric, long-term benefits and risks. *Future Sci OA* 2(2):FSO116.
4. MRC-Vitamin-Study-Research-Group (1991) Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. *Lancet* 338(8760):131-137.
5. Czeizel A & Dudas I (1992) Prevention of the first occurrence of neural tube defects by periconceptional vitamin supplementation. *N Eng J Med* 327(26):1832-1835.
6. Smithells RW, Sheppard S, Wild J, & Schorah CJ (1989) Prevention of neural tube defect recurrences in Yorkshire: final report. *Lancet* 2(8661):498-499.
7. Nevin NC & Seller MJ (1990) Prevention of neural-tube-defect recurrences. *Lancet* 335(8682):178-179.
8. Standing-Committee-on-the-Scientific-Evaluation-of-Dietary-Reference-Intakes (1998) Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline, The National Academies Collection: Reports funded by National Institutes of Health, Washington (DC)).
9. Centers-for-Disease-Control (1992) Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. in *MMWR Recomm Rep*, pp 1-7.
10. Finer LB & Henshaw SK (2006) Disparities in rates of unintended pregnancy in the United States, 1994 and 2001. *Perspectives on sexual and reproductive health* 38(2):90-96.
11. Crozier SR, et al. (2009) Do women change their health behaviours in pregnancy? Findings from the Southampton Women's Survey. *Paediatr Perinatal Epidemiol* 23(5):446-453.
12. Food-Fortification-Initiative (2016) Say Hello to a Fortified Future, 2016 in Review. (Food Fortification Initiative, Atlanta, USA).
13. Searby L (2016) Folic acid fortification: The current global state of play.
14. Hopkins SM, et al. (2015) Impact of voluntary fortification and supplement use on dietary intakes and biomarker status of folate and vitamin B-12 in Irish adults. *Am J Clin Nutr* 101(6):1163-1172.
15. Daly LE, Kirke PN, Molloy A, Weir DG, & Scott JM (1995) Folate levels and neural tube defects. Implications for prevention. *JAMA* 274(21):1698-1702.

16. Crider KS, et al. (2014) Population red blood cell folate concentrations for prevention of neural tube defects: Bayesian model. *B Med J* 349:g4554.
17. Food-and-Drug-Administration-HHS (1993) Food standards: amendment of the standards of identity for enriched grain products to require addition of folic acid. 58(197):53305-53312.
18. Bailey LB (2004) Folate and vitamin B12 recommended intakes and status in the United States. *Nutr Rev* 62(6 Pt 2):S14-20; discussion S21.
19. Yang Q, et al. (2010) Folic acid source, usual intake, and folate and vitamin B-12 status in US adults: National Health and Nutrition Examination Survey (NHANES) 2003-2006. *Am J Clin Nutr* 91(1):64-72.
20. Carmel R & Johnson CS (1978) Racial patterns in pernicious anemia. Early age at onset and increased frequency of intrinsic-factor antibody in black women. *New Engl J Med* 298(12):647-650.
21. Hannibal L, et al. (2016) Biomarkers and Algorithms for the Diagnosis of Vitamin B12 Deficiency. *Front Mol Biosci* 3:27.
22. Kriebel D & Tickner J (2001) Reenergizing Public Health Through Precaution. *Am J Publ Health* 91(9):1351-1355.
23. Goldstein BD (2001) The precautionary principle also applies to public health actions. *Am J Public Health* 91(9):1358-1361.
24. World-Health-Organization (2015) Optimal serum and red blood cell folate concentrations in women of reproductive age for prevention of neural tube defects.
25. Yetley EA, et al. (2017) Options for basing Dietary Reference Intakes (DRIs) on chronic disease endpoints: report from a joint US-/Canadian-sponsored working group. *Am J Clin Nutr* 105(1):249S-285S.
26. Detrait ER, et al. (2005) Human neural tube defects: developmental biology, epidemiology, and genetics. *Neurotox Teratol* 27(3):515-524.
27. Gelineau-van Waes J, et al. (2012) Increased sphingoid base-1-phosphates and failure of neural tube closure after exposure to fumonisin or FTY720. *Birth defects research. Part A. Clin Mol Teratol* 94(10):790-803.
28. Mazumdar M, et al. (2015) Arsenic is associated with reduced effect of folic acid in myelomeningocele prevention: a case control study in Bangladesh. *Environ Health* 14:34.
29. Suh JR, Herbig AK, & Stover PJ (2001) New perspectives on folate catabolism. *Ann Rev Nutr* 21:255-282.
30. Suitor CW & Bailey LB (2000) Dietary folate equivalents: interpretation and application. *J Am Diet Assoc* 100(1):88-94.
31. Bailey SW & Ayling JE (2009) The extremely slow and variable activity of dihydrofolate reductase in human liver and its implications for high folic acid intake. *Proc Nat Acad Sci* 106(36):15424-15429.

32. Pfeiffer CM, et al. (2015) Unmetabolized folic acid is detected in nearly all serum samples from US children, adolescents, and adults. *J Nutr* 145(3):520-531.
33. Wright AJ, et al. (2005) Differential kinetic behavior and distribution for pteroylglutamic acid and reduced folates: a revised hypothesis of the primary site of PteGlu metabolism in humans. *J Nutr* 135(3):619-623.
34. West AA, et al. (2012) Folate-status response to a controlled folate intake in nonpregnant, pregnant, and lactating women. *Am J Clin Nutr* 96(4):789-800.
35. Carmel R (2008) Nutritional anemias and the elderly. *Semin Hematol* 45(4):225-234.
36. Agamanolis DP, et al. (1976) Neuropathology of experimental vitamin B12 deficiency in monkeys. *Neurology* 26(10):905-914.
37. Yeung L, Yang Q, & Berry RJ (2008) Contributions of total daily intake of folic acid to serum folate concentrations. *JAMA* 300(21):2486-2487.
38. Scientific-Acvisory-Committee-on-Nutrition (2017) Draft--Update on folic acid.
39. Troen AM, et al. (2006) Unmetabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J Nutr* 136(1):189-194.
40. Kelly P, McPartlin J, Goggins M, Weir DG, & Scott JM (1997) Unmetabolized folic acid in serum: acute studies in subjects consuming fortified food and supplements. *Am J Clin Nutr* 65(6):1790-1795.
41. Pfeiffer CM, et al. (2015) Folate status and concentrations of serum folate forms in the US population: National Health and Nutrition Examination Survey 2011-2. *Br J Nutr* 113(12):1965-1977.
42. Fox JT & Stover PJ (2008) Folate-mediated one-carbon metabolism. *Vitamins and hormones* 79:1-44.
43. Yetley EA, et al. (2011) Biomarkers of folate status in NHANES: a roundtable summary. *Am J Clin Nutr* 94(1):303S-312S.
44. Misselbeck K, et al. (2017) A hybrid stochastic model of folate-mediated one-carbon metabolism: Effect of the common C677T MTHFR variant on de novo thymidylate biosynthesis. *Sci Rep* 7(1):797.
45. Boyles AL, Yetley EA, Thayer KA, & Coates PM (2016) Safe use of high intakes of folic acid: research challenges and paths forward. *Nutr Rev* 74(7):469-474.
46. National-Toxicology-Program (2015) NTP Monograph: Identifying Research Needs for Assessing Safe Use of High Intakes of Folic Acid.
47. Food-Safety-Authority-of-Ireland (2016) Update Report on Folic Acid and the Prevention of Birth Defects in Ireland. (Dublin, Ireland).
48. Food-and-Drug-Administration-HHS (2016) Food Additives Permitted for Direct Addition to Food for Human Consumption; Folic Acid.

49. Norwegian-Scientific-Committee-for-Food-Safety (2015) Risk assessment of folic acid and food supplements; opinion of the panel on nutrition, dietetic products, novel food and allergy of the Norwegian Scientific Committee for Food Safety (VKM). (Oslo, Norway).
50. European-Food-Safety-Authority (2014) Update Report on Folic Acid and the Prevention of Birth Defects in Ireland.
51. Smith AD, Kim YI, & Refsum H (2008) Is folic acid good for everyone? *Am J Clin Nutr* 87(3):517-533.
52. Mason JB (2009) Folate, cancer risk, and the Greek god, Proteus: a tale of two chameleons. *Nutr Rev* 67(4):206-212.
53. Kim YI (2006) Folate: a magic bullet or a double edged sword for colorectal cancer prevention? *Gut* 55(10):1387-1389.
54. Vollset SE, et al. (2013) Effects of folic acid supplementation on overall and site-specific cancer incidence during the randomised trials: meta-analyses of data on 50,000 individuals. *Lancet* 381(9871):1029-1036.
55. Qin T, et al. (2015) Folic acid supplements and colorectal cancer risk: meta-analysis of randomized controlled trials. *Sci Rep* 5:12044.
56. Mason JB (2011) Folate consumption and cancer risk: a confirmation and some reassurance, but we're not out of the woods quite yet. *Am J Clin Nutr* 94(4):965-966.
57. Teh AH, Symonds E, Bull C, Clifton P, & Fenech M (2012) The influence of folate and methionine on intestinal tumour development in the Apc(Min/+) mouse model. *Mut Res*.
58. Palmer AM, Kamynina E, Field MS, & Stover PJ (2017) Folate rescues vitamin B12 depletion-induced inhibition of nuclear thymidylate biosynthesis and genome instability. *Proc Nat Acad Sci*.
59. Lindenbaum J, et al. (1995) Neuropsychiatric disorders caused by cobalamin deficiency in the absence of anemia or macrocytosis. 1988. *Nutrition* 11(2):181; discussion 180, 182.
60. Morris MS, Jacques PF, Rosenberg IH, & Selhub J (2007) Folate and vitamin B-12 status in relation to anemia, macrocytosis, and cognitive impairment in older Americans in the age of folic acid fortification. *Am J Clin Nutr* 85(1):193-200.
61. Morris MS, Selhub J, & Jacques PF (2012) Vitamin B-12 and Folate Status in Relation to Decline in Scores on the Mini-Mental State Examination in the Framingham Heart Study. *J Am Ger Soc* 60(8):1457-1464.
62. Miller JW, Garrod MG, Allen LH, Haan MN, & Green R (2009) Metabolic evidence of vitamin B-12 deficiency, including high homocysteine and methylmalonic acid and low holotranscobalamin, is more pronounced in older adults with elevated plasma folate. *Am J Clin Nutr* 90(6):1586-1592.
63. Young SS (2011) Deming, data and observational studies. *Significance*:116-120.

64. Brito A, et al. (2016) Vitamin B-12 treatment of asymptomatic, deficient, elderly Chileans improves conductivity in myelinated peripheral nerves, but high serum folate impairs vitamin B-12 status response assessed by the combined indicator of vitamin B-12 status. *Am J Clin Nutr* 103(1):250-257.
65. Yajnik CS, et al. (2008) Vitamin B12 and folate concentrations during pregnancy and insulin resistance in the offspring: the Pune Maternal Nutrition Study. *Diabetologia* 51(1):29-38.
66. Stewart CP, et al. (2009) Antenatal micronutrient supplementation reduces metabolic syndrome in 6- to 8-year-old children in rural Nepal. *J Nutr* 139(8):1575-1581.
67. Cooney CA, Dave AA, & Wolff GL (2002) Maternal methyl supplements in mice affect epigenetic variation and DNA methylation of offspring. *J Nutr* 132(8 Suppl):2393S-2400S.
68. Dolinoy DC, Weidman JR, Waterland RA, & Jirtle RL (2006) Maternal genistein alters coat color and protects Avy mouse offspring from obesity by modifying the fetal epigenome. *Environ Health Pers* 114(4):567-572.
69. Morgan HD, Sutherland HGE, Martin DIK, & Whitelaw E (1999) Epigenetic inheritance at the agouti locus in the mouse. *Nat Genet* 23(3):314-318.
70. Waterland RA & Jirtle RL (2003) Transposable elements: targets for early nutritional effects on epigenetic gene regulation. *Mol Cell Biol* 23(15):5293-5300.
71. Modabbernia A, Velthorst E, & Reichenberg A (2017) Environmental risk factors for autism: an evidence-based review of systematic reviews and meta-analyses. *Molecular autism* 8:13.
72. Gao Y, et al. (2016) New Perspective on Impact of Folic Acid Supplementation during Pregnancy on Neurodevelopment/Autism in the Offspring Children – A Systematic Review. *PLOS ONE* 11(11):e0165626.
73. Zhao M, et al. (2017) Homocysteine and Stroke Risk: Modifying Effect of Methylenetetrahydrofolate Reductase C677T Polymorphism and Folic Acid Intervention. *Stroke*.
74. Xu A, et al. (2016) A Meta-Analysis of the Relationship Between Maternal Folic Acid Supplementation and the Risk of Congenital Heart Defects. *Int Heart J* 57(6):725-728.
75. Qin X, et al. (2016) Folic Acid Therapy Reduces the First Stroke Risk Associated With Hypercholesterolemia Among Hypertensive Patients. *Stroke* 47(11):2805-2812.
76. Liu S, et al. (2016) Effect of Folic Acid Food Fortification in Canada on Congenital Heart Disease Subtypes. *Circulation* 134(9):647-655.

SHOULD VITAMIN B12 BE CONSIDERED IN ASSESSING NTD RISK

1. Blencowe, H., et al., Folic acid to reduce neonatal mortality from neural tube disorders. *Int J Epidemiol*, 2010. 39 Suppl 1: p. i110-21.
2. Matthews, R.G. Cobalamin-dependent methionine synthase, in *Chemistry and Biochemistry of B12*, R. Banerjee, Editor. 1999, John Wiley and Sons, Inc: New York. p. 681-706.
3. Scott, J.M. and D.G. Weir. The methyl folate trap. A physiological response in man to prevent methyl group deficiency in kwashiorkor (methionine deficiency) and an explanation for folic-acid induced exacerbation of subacute combined degeneration in pernicious anaemia. *Lancet*, 1981. 2(8242): p. 337-40.
4. Smithells, R.W., S. Sheppard, and C.J. Schorah. Vitamin deficiencies and neural tube defects. *Arch Dis Child*, 1976. 51(12): p. 944-50.
5. Smithells, R.W., et al. Further experience of vitamin supplementation for prevention of neural tube defect recurrences. *Lancet*, 1983. 1(8332): p. 1027-31.
6. Schorah, C.J., et al. The effect of periconceptional supplementation on blood vitamin concentrations in women at recurrence risk for neural tube defect. *Br J Nutr*, 1983. 49(2): p. 203-11.
7. MRC. Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. MRC Vitamin Study Research Group. *Lancet*, 1991. 338(8760): p. 131-7.
8. Molloy, A.M., et al. Effects of folate and vitamin B12 deficiencies during pregnancy on fetal, infant, and child development. *Food Nutr Bull*, 2008. 29(2 Suppl): p. S101-11; discussion S112-5.
9. Allen, L.H. How common is vitamin B-12 deficiency? *Am J Clin Nutr*, 2009. 89(2): p. 693s-6s.
10. Schorah, C.J., R.W. Smithells, and J. Scott. Vitamin B12 and anencephaly. *Lancet*, 1980. 1(8173): p. 880.
11. Steegers-Theunissen, R.P., et al. Neural-tube defects and derangement of homocysteine metabolism. *N Engl J Med*, 1991. 324(3): p. 199-200.
12. Molloy, A.M., et al. Maternal serum folate and vitamin-B12 concentrations in pregnancies associated with neural-tube defects. *Arch Dis Child*, 1985. 60(7): p. 660-665.
13. Yates, J.R., et al. Is disordered folate metabolism the basis for the genetic predisposition to neural tube defects? *Clin Genet*, 1987. 31(5): p. 279-87.
14. Afman, L.A., et al. Reduced vitamin B12 binding by transcobalamin II increases the risk of neural tube defects. *QJM*, 2001. 94(3): p. 159-66.
15. Wild, J., et al. Investigation of factors influencing folate status in women who have had a neural tube defect-affected infant. *Br J Obstet Gynaecol*, 1993. 100(6): p. 546-9.

16. Wald, N.J., et al. Blood folic acid and vitamin B12 in relation to neural tube defects. *Br J Obstet Gynaecol*, 1996. 103(4): p. 319-24.
17. Steegers-Theunissen, R.P., et al. Neural tube defects and elevated homocysteine levels in amniotic fluid. *Am J Obstet Gynecol*, 1995. 172(5): p. 1436-41.
18. Mills, J.L., et al. Homocysteine metabolism in pregnancies complicated by neural-tube defects. *Lancet*, 1995. 345(8943): p. 149-51.
19. Gardiki-Kouidou, P. and M.J. Seller. Amniotic fluid folate, vitamin B12 and transcobalamins in neural tube defects. *Clin Genet*, 1988. 33(6): p. 441-8.
20. Dawson, E.B., D.R. Evans, and J.W. Van Hook. Amniotic fluid B12 and folate levels associated with neural tube defects. *Am J Perinatol*, 1998. 15(9): p. 511-4.
21. Steen, M.T., et al. Neural-tube defects are associated with low concentrations of cobalamin (vitamin B12) in amniotic fluid. *Prenat Diagn*, 1998. 18(6): p. 545-55.
22. Weekes, E.W., et al. Nutrient levels in amniotic fluid from women with normal and neural tube defect pregnancies. *Biol Neonate*, 1992. 61(4): p. 226-31.
23. Economides, D.L., et al. Folate and vitamin B12 concentrations in maternal and fetal blood, and amniotic fluid in second trimester pregnancies complicated by neural tube defects. *Br J Obstet Gynaecol*, 1992. 99(1): p. 23-5.
24. Mills, J.L., et al. Maternal vitamin levels during pregnancies producing infants with neural tube defects. *J Pediatr*, 1992. 120(6): p. 863-71.
25. Molloy, A.M., et al. Maternal vitamin B12 status and risk of neural tube defects in a population with high neural tube defect prevalence and no folic acid fortification. *Pediatrics*, 2009. 123(3): p. 917-923.
26. Zhang, T., et al. Maternal serum vitamin B12, folate and homocysteine and the risk of neural tube defects in the offspring in a high-risk area of China. *Public Health Nutr*, 2009. 12(5): p. 680-6.
27. Suarez, L., et al. Maternal serum B12 levels and risk for neural tube defects in a Texas-Mexico border population. *Ann Epidemiol*, 2003. 13(2): p. 81-8.
28. Ray, J.G., et al. Vitamin B12 and the risk of neural tube defects in a folic-acid-fortified population. *Epidemiology*, 2007. 18(3): p. 362-6.
29. Nasri, K., et al. Association of maternal homocysteine and vitamins status with the risk of neural tube defects in Tunisia: A case-control study. *Birth Defects Res A Clin Mol Teratol*, 2015. 103(12): p. 1011-20.
30. Tang, K.F., Y.L. Li, and H.Y. Wang. Quantitative assessment of maternal biomarkers related to one-carbon metabolism and neural tube defects. *Sci Rep*, 2015. 5: p. 8510.
31. Peker, E., et al. The levels of vitamin B12, folate and homocysteine in mothers and their babies with neural tube defects. *J Matern Fetal Neonatal Med*, 2016. 29(18): p. 2944-8.

32. Czeizel, A.E. and I. Dudas. Prevention of the first occurrence of neural-tube defects by periconceptual vitamin supplementation. *N Engl J Med*, 1992. 327(26): p. 1832-5.
33. Molloy, A.M., et al. The search for genetic polymorphisms in the homocysteine/folate pathway that contribute to the etiology of human neural tube defects. *Birth Defects Res A Clin Mol Teratol*, 2009. 85(4): p. 285-94.
34. Amorim, M.R., et al. Non-Latin European descent could be a requirement for association of NTD and MTHFR variant 677C > T: a meta-analysis. *Am J Med Genet A*, 2007. 143A(15): p. 1726-32.
35. Shields, D.C., et al. The "thermolabile" variant of methylenetetrahydrofolate reductase and neural tube defects: An evaluation of genetic risk and the relative importance of the genotypes of the embryo and the mother. *Am J Hum Genet*, 1999. 64(4): p. 1045-55.
36. Boyles, A.L., et al. Neural tube defects and folate pathway genes: family-based association tests of gene-gene and gene-environment interactions. *Environ Health Perspect*, 2006. 114(10): p. 1547-52.
37. Pangilinan, F., et al. Transcobalamin II polymorphisms influence plasma levels of vitamin B12 but are not major risk factors for neural tube defects. *American Journal of Human Genetics*, 2003. 73(5): p. 525-525.
38. Candito, M., et al. Nutritional and genetic determinants of vitamin B and homocysteine metabolisms in neural tube defects: A multicenter case-control study. *Am J Med Genet A*, 2008. 146A(9): p. 1128-1133.
39. Pangilinan, F., et al. Replication and exploratory analysis of 24 candidate risk polymorphisms for neural tube defects. *BMC Med Genet*, 2014. 15(1): p. 102.
40. Winkels, R.M., et al. Bread cofortified with folic acid and vitamin B-12 improves the folate and vitamin B-12 status of healthy older people: a randomized controlled trial. *Am J Clin Nutr*, 2008. 88(2): p. 348-55.
41. Swanson, D.A., et al. Evaluation of transcobalamin II polymorphisms as neural tube defect risk factors in an Irish population. *Birth Defects Res A Clin Mol Teratol*, 2005. 73(4): p. 239-44.
42. Afman, L.A., et al. Single nucleotide polymorphisms in the transcobalamin gene: relationship with transcobalamin concentrations and risk for neural tube defects. *Eur J Hum Genet*, 2002. 10(7): p. 433-8.
43. Pangilinan, F., et al. Evaluation of common genetic variants in 82 candidate genes as risk factors for neural tube defects. *BMC Med Genet*, 2012. 13: p. 62.
44. Pangilinan, F., et al. Transcobalamin II receptor polymorphisms are associated with increased risk for neural tube defects. *Journal of Medical Genetics*, 2010. 47(10): p. 677-685.

45. Kirke, P.N., et al. Maternal plasma folate and vitamin B12 are independent risk factors for neural tube defects. *Q J Med*, 1993. 86(11): p. 703-8.
46. Afman, L.A., et al. Gene-gene interaction between the cystathionine beta-synthase 31 base pair variable number of tandem repeats and the methylenetetrahydrofolate reductase 677C > T polymorphism on homocysteine levels and risk for neural tube defects. *Mol Genet Metab*, 2003. 78(3): p. 211-5.
47. Bjorke-Monsen, A.L., et al. Elevated plasma total homocysteine and C677T mutation of the methylenetetrahydrofolate reductase gene in patients with spina bifida. *Qjm*, 1997. 90(9): p. 593-6.
48. Daly, L.E., et al. Folate levels and neural tube defects. Implications for prevention. *JAMA*, 1995. 274(21): p. 1698-702.
49. Crider, K.S., et al. Population red blood cell folate concentrations for prevention of neural tube defects: Bayesian model. *Bmj*, 2014. 349: p. g4554.
50. Rosenthal, J., et al. Folate Deficiency Is Prevalent in Women of Childbearing Age in Belize and Is Negatively Affected by Coexisting Vitamin B-12 Deficiency: Belize National Micronutrient Survey 2011. *J Nutr*, 2017.
51. Bailey, R.L., et al. Modeling a methylmalonic acid-derived change point for serum vitamin B-12 for adults in NHANES. *Am J Clin Nutr*, 2013. 98(2): p. 460-7.
52. Metz, J. Haematological implications of folate food fortification. *S Afr Med J*, 2013. 103(12 Suppl 1): p. 978-81.
53. Hazra, A., et al. Common variants of FUT2 are associated with plasma vitamin B12 levels. *Nat Genet*, 2008. 40(10): p. 1160-2.
54. Tanaka, T., et al. Genome-wide association study of vitamin B6, vitamin B12, folate, and homocysteine blood concentrations. *Am J Hum Genet*, 2009. 84(4): p. 477-82.
55. Murphy, M.M., et al. Longitudinal study of the effect of pregnancy on maternal and fetal cobalamin status in healthy women and their offspring. *J Nutr*, 2007. 137(8): p. 1863-7.
56. Benson, J., et al. Low vitamin B12 levels among newly-arrived refugees from Bhutan, Iran and Afghanistan: a multicentre Australian study. *PLoS One*, 2013. 8(2): p. e57998.
57. Calis, J.C., et al. Severe anemia in Malawian children. *N Engl J Med*, 2008. 358(9): p. 888-99.
58. Dang, S., et al. The status of vitamin B12 and folate among Chinese women: a population-based cross-sectional study in northwest China. *PLoS One*, 2014. 9(11): p. e112586.
59. Margo, G., et al. Protein energy malnutrition and nutritional anaemia in preschool children in rural KwaZulu. *S Afr Med J*, 1978. 53(1): p. 21-6.
60. McLean, E., B. de Benoist, and L.H. Allen. Review of the magnitude of folate and vitamin B12 deficiencies worldwide. *Food Nutr Bull*, 2008. 29(2 Suppl): p. S38-51.

61. Pawlak, R., S.E. Lester, and T. Babatunde. The prevalence of cobalamin deficiency among vegetarians assessed by serum vitamin B12: a review of literature. *Eur J Clin Nutr*, 2014. 68(5): p. 541-8.
62. Yetley, E.A., et al. Biomarkers of vitamin B-12 status in NHANES: a roundtable summary. *Am J Clin Nutr*, 2011. 94(1): p. 313S-321S.
63. Sobczyńska-Malefora, A., et al. An audit of holotranscobalamin ("Active" B 12) and methylmalonic acid assays for the assessment of vitamin B 12 status: Application in a mixed patient population. *Clinical biochemistry*, 2014. 47(1): p. 82-86.
64. Allen, L.H., et al. Vitamin B-12 deficiency and malabsorption are highly prevalent in rural Mexican communities. *Am J Clin Nutr*, 1995. 62(5): p. 1013-9.
65. Bailey, R.L., et al. Monitoring of vitamin B-12 nutritional status in the United States by using plasma methylmalonic acid and serum vitamin B-12. *Am J Clin Nutr*, 2011. 94(2): p. 552-61.
66. Refsum, H., et al. Hyperhomocysteinemia and elevated methylmalonic acid indicate a high prevalence of cobalamin deficiency in Asian Indians. *Am J Clin Nutr*, 2001. 74(2): p. 233-41.
67. Bondevik, G.T., et al. Homocysteine and methylmalonic acid levels in pregnant Nepali women. Should cobalamin supplementation be considered? *Eur J Clin Nutr*, 2001. 55(10): p. 856-64.
68. Fedosov, S.N., et al. Combined indicator of vitamin B12 status: modification for missing biomarkers and folate status and recommendations for revised cut-points. *Clin Chem Lab Med*, 2015.
69. van der Westhuyzen, J., et al. Iron, folate and vitamin B12 nutrition and anaemia in black preschool children in the northern Transvaal. *S Afr Med J*, 1986. 70(3): p. 143-6.
70. Bhardwaj, A., et al. Rapid Assessment for Coexistence of Vitamin B12 and Iron Deficiency Anemia among Adolescent Males and Females in Northern Himalayan State of India. *Anemia*, 2013. 2013: p. 959605.
71. Pathak, P., et al. Iron, folate, and vitamin B12 stores among pregnant women in a rural area of Haryana State, India. *Food Nutr Bull*, 2007. 28(4): p. 435-8.
72. Devalia, V., M.S. Hamilton, and A.M. Molloy. Guidelines for the diagnosis and treatment of cobalamin and folate disorders. *Br J Haematol*, 2014.
73. Carmel, R. and M. Sarrai. Diagnosis and management of clinical and subclinical cobalamin deficiency: advances and controversies. *Curr Hematol Rep*, 2006. 5(1): p. 23-33.
74. Carmel, R. Diagnosis and management of clinical and subclinical cobalamin deficiencies: why controversies persist in the age of sensitive metabolic testing. *Biochimie*, 2013. 95(5): p. 1047-55.

75. Mosen, A.L., et al. Cobalamin status and its biochemical markers methylmalonic acid and homocysteine in different age groups from 4 days to 19 years. *Clin Chem*, 2003. 49(12): p. 2067-75.
76. Ray, J.G., et al. High rate of maternal vitamin B12 deficiency nearly a decade after Canadian folic acid flour fortification. *QJM*, 2008.
77. Hopkins, S.M., et al. Impact of voluntary fortification and supplement use on dietary intakes and biomarker status of folate and vitamin B-12 in Irish adults. *The American journal of clinical nutrition*, 2015: p. ajcn107151.
78. Brito, A., et al. Folate and Vitamin B12 Status in Latin America and the Caribbean: An Update. *Food Nutr Bull*, 2015. 36(2 Suppl): p. S109-18.
79. Garcia-Casal, M.N., et al. High prevalence of folic acid and vitamin B12 deficiencies in infants, children, adolescents and pregnant women in Venezuela. *Eur J Clin Nutr*, 2005. 59(9): p. 1064-70.
80. Vitamin B12 deficiency in resettled Bhutanese refugees--United States, 2008-2011. *MMWR Morb Mortal Wkly Rep*, 2011. 60(11): p. 343-6.
81. Yajnik, C.S., et al. Oral vitamin B12 supplementation reduces plasma total homocysteine concentration in women in India. *Asia Pac J Clin Nutr*, 2007. 16(1): p. 103-9.
82. Yajnik, C.S., et al. Maternal total homocysteine concentration and neonatal size in India. *Asia Pac J Clin Nutr*, 2005. 14(2): p. 179-81.
83. Jiang, T., et al. Micronutrient deficiencies in early pregnancy are common, concurrent, and vary by season among rural Nepali pregnant women. *J Nutr*, 2005. 135(5): p. 1106-12.
84. Kapil, U. and A.S. Bhadoria. Prevalence of Folate, Ferritin and Cobalamin Deficiencies amongst Adolescent in India. *J Family Med Prim Care*, 2014. 3(3): p. 247-9.
85. Godbole, K., U. Deshmukh, and C. Yajnik. Nutrigenetic determinants of neural tube defects in India. *Indian Pediatr*, 2009. 46(6): p. 467-75.
86. Neumann, C.G., et al. Low vitamin B12 intake during pregnancy and lactation and low breastmilk vitamin 12 content in rural Kenyan women consuming predominantly maize diets. *Food Nutr Bull*, 2013. 34(2): p. 151-9.
87. Shahab-Ferdows, S., et al. Regional, Socioeconomic, and Dietary Risk Factors for Vitamin B-12 Deficiency Differ from Those for Folate Deficiency in Cameroonian Women and Children. *J Nutr*, 2015. 145(11): p. 2587-95.
88. Wirth, J.P., et al. Anemia, Micronutrient Deficiencies, and Malaria in Children and Women in Sierra Leone Prior to the Ebola Outbreak - Findings of a Cross-Sectional Study. *PLoS One*, 2016. 11(5): p. e0155031.

89. Harvey-Leeson, S., et al. Anemia and Micronutrient Status of Women of Childbearing Age and Children 6-59 Months in the Democratic Republic of the Congo. *Nutrients*, 2016. 8(2): p. 98.
90. Dei-Adomakoh, Y., et al. Second Trimester Anaemia in Pregnant Ghanaians. *West Afr J Med*, 2014. 33(4): p. 229-33.
91. Fernandes-Costa, F. and J. Metz. A comparison of serum transcobalamin levels in white and black subjects. *Am J Clin Nutr*, 1982. 35(1): p. 83-6.
92. Fleming, A.F., Y.A. Ogunfunmilade, and R. Carmel, Serum vitamin B12 levels and vitamin B12-binding proteins of serum and saliva of healthy Nigerians and Europeans. *Am J Clin Nutr*, 1978. 31(10): p. 1732-8.
93. Saxena, S. and R. Carmel. Racial differences in vitamin B12 levels in the United States. *Am J Clin Pathol*, 1987. 88(1): p. 95-7.
94. Stabler, S.P., et al. Racial differences in prevalence of cobalamin and folate deficiencies in disabled elderly women. *Am J Clin Nutr*, 1999. 70(5): p. 911-9.
95. Pfeiffer, C.M., et al. Trends in blood folate and vitamin B-12 concentrations in the United States, 1988 2004. *Am J Clin Nutr*, 2007. 86(3): p. 718-27.
96. Oussalah, A., et al. Helicobacter pylori serologic status has no influence on the association between fucosyltransferase 2 polymorphism (FUT2 461 G->A) and vitamin B-12 in Europe and West Africa. *Am J Clin Nutr*, 2012. 95(2): p. 514-21.
97. Kelly, R.J., et al. Sequence and expression of a candidate for the human Secretor blood group alpha(1,2)fucosyltransferase gene (FUT2). Homozygosity for an enzyme-inactivating nonsense mutation commonly correlates with the non-secretor phenotype. *J Biol Chem*, 1995. 270(9): p. 4640-9.
98. Marionneau, S., et al. ABH and Lewis histo-blood group antigens, a model for the meaning of oligosaccharide diversity in the face of a changing world. *Biochimie*, 2001. 83(7): p. 565-73.
99. Segurel, L., et al. The ABO blood group is a trans-species polymorphism in primates. *Proc Natl Acad Sci U S A*, 2012. 109(45): p. 18493-8.
100. Molloy, A.M., et al. Low blood folates in NTD pregnancies are only partly explained by thermolabile 5,10-methylenetetrahydrofolate reductase: low folate status alone may be the critical factor. *Am J Med Genet*, 1998. 78(2): p. 155-9.
101. Molloy, A.M., et al. Thermolabile variant of 5,10-methylenetetrahydrofolate reductase associated with low red-cell folates: implications for folate intake recommendations. *Lancet*, 1997. 349(9065): p. 1591-3.
102. Yang, Q.H., et al. Prevalence and effects of gene-gene and gene-nutrient interactions on serum folate and serum total homocysteine concentrations in the United States: findings

- from the third National Health and Nutrition Examination Survey DNA Bank. *Am J Clin Nutr*, 2008. 88(1): p. 232-46.
103. Wilcken, B., et al. Geographical and ethnic variation of the 677C>T allele of 5,10 methylenetetrahydrofolate reductase (MTHFR): findings from over 7000 newborns from 16 areas world wide. *J Med Genet*, 2003. 40(8): p. 619-25.
 104. Botto, L.D. and Q. Yang. 5,10-Methylenetetrahydrofolate reductase gene variants and congenital anomalies: a HuGE review. *Am J Epidemiol*, 2000. 151(9): p. 862-77.
 105. Mutchinick, O.M., et al. High prevalence of the thermolabile methylenetetrahydrofolate reductase variant in Mexico: a country with a very high prevalence of neural tube defects. *Mol Genet Metab*, 1999. 68(4): p. 461-7.
 106. Crider, K.S., et al. MTHFR 677C->T genotype is associated with folate and homocysteine concentrations in a large, population-based, double-blind trial of folic acid supplementation. *Am J Clin Nutr*, 2011. 93(6): p. 1365-72.
 107. Williams, L.J., et al. Decline in the prevalence of spina bifida and anencephaly by race/ethnicity: 1995-2002. *Pediatrics*, 2005. 116(3): p. 580-6.
 108. Zaganjor, I., et al. Describing the Prevalence of Neural Tube Defects Worldwide: A Systematic Literature Review. *PLoS One*, 2016. 11(4): p. e0151586.
 109. Besser, L.M., L.J. Williams, and J.D. Cragan. Interpreting changes in the epidemiology of anencephaly and spina bifida following folic acid fortification of the U.S. grain supply in the setting of long-term trends, Atlanta, Georgia, 1968-2003. *Birth Defects Res A Clin Mol Teratol*, 2007. 79(11): p. 730-6.
 110. Chanarin, I. *The Megaloblastic Anaemias*. 1st Edition ed. 1969, Oxford and Edinburgh: Blackwell Scientific Publications.
 111. Bor, M.V., et al. A daily intake of approximately 6 microg vitamin B-12 appears to saturate all the vitamin B-12-related variables in Danish postmenopausal women. *Am J Clin Nutr*, 2006. 83(1): p. 52-8.
 112. Abdeen, Z., et al. Predicted efficacy of the Palestinian wheat flour fortification programme: complementary analysis of biochemical and dietary data. *Public Health Nutr*, 2015. 18(8): p. 1358-68.
 113. Dullemeijer, C., et al. Systematic review with dose-response meta-analyses between vitamin B-12 intake and European Micronutrient Recommendations Aligned's prioritized biomarkers of vitamin B-12 including randomized controlled trials and observational studies in adults and elderly persons. *Am J Clin Nutr*, 2013. 97(2): p. 390-402.
 114. Dangour, A.D., et al. A randomised controlled trial investigating the effect of vitamin B12 supplementation on neurological function in healthy older people: the Older People and Enhanced Neurological function (OPEN) study protocol [ISRCTN54195799]. *Nutr J*, 2011. 10: p. 22.

115. Katre, P., et al. Vitamin B12 and folic acid supplementation and plasma total homocysteine concentrations in pregnant Indian women with low B12 and high folate status. *Asia Pac J Clin Nutr*, 2010. 19(3): p. 335-43.
116. Hill, M.H., et al. A vitamin B-12 supplement of 500 mug/d for eight weeks does not normalize urinary methylmalonic acid or other biomarkers of vitamin B-12 status in elderly people with moderately poor vitamin B-12 status. *J Nutr*, 2013. 143(2): p. 142-7.
117. Deshmukh, U.S., et al. Effect of physiological doses of oral vitamin B12 on plasma homocysteine: a randomized, placebo-controlled, double-blind trial in India. *Eur J Clin Nutr*, 2010. 64(5): p. 495-502.
118. Eussen, S.J., et al. Oral cyanocobalamin supplementation in older people with vitamin B12 deficiency: a dose-finding trial. *Arch Intern Med*, 2005. 165(10): p. 1167-72.
119. Allen, L.H., et al. Considering the case for vitamin B12 fortification of flour. *Food Nutr Bull*, 2010. 31(1 Suppl): p. S36-46.
120. Taneja, S., et al. Folic acid and vitamin B-12 supplementation and common infections in 6-30-mo-old children in India: a randomized placebo-controlled trial. *Am J Clin Nutr*, 2013.
121. Carmel, R.. Mandatory fortification of the food supply with cobalamin: an idea whose time has not yet come. *J Inherit Metab Dis*, 2011. 34(1): p. 67-73.
122. Refsum, H. and A.D. Smith. Are we ready for mandatory fortification with vitamin B-12? *Am J Clin Nutr*, 2008. 88(2): p. 253-4.
123. Carmel, R. Efficacy and safety of fortification and supplementation with vitamin B12: biochemical and physiological effects. *Food Nutr Bull*, 2008. 29(2 Suppl): p. S177-87.
124. Bae, S., et al. Vitamin B-12 Status Differs among Pregnant, Lactating, and Control Women with Equivalent Nutrient Intakes. *J Nutr*, 2015. 145(7): p. 1507-14.
125. Duggan, C., et al. Vitamin B-12 Supplementation during Pregnancy and Early Lactation Increases Maternal, Breast Milk, and Infant Measures of Vitamin B-12 Status. *J Nutr*, 2014.
126. Yajnik, C.S., et al. Vitamin B12 and folate concentrations during pregnancy and insulin resistance in the offspring: the Pune Maternal Nutrition Study. *Diabetologia*, 2008. 51(1): p. 29-38.
127. Institute of Medicine Standing Committee on the Scientific Evaluation of Dietary Reference, Intake, Panel on Folate, and Choline. The National Academies Collection: Reports funded by National Institutes of Health, in Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline. 1998, National Academies Press (US) National Academy of Sciences.: Washington (DC).
128. Allen, R.H. and S.P. Stabler Identification and quantitation of cobalamin and cobalamin analogues in human feces. *Am J Clin Nutr*, 2008. 87(5): p. 1324-35.

FRAMEWORK FOR LABORATORY HARMONIZATION OF FOLATE MEASUREMENTS IN LOW-AND-MIDDLE-INCOME COUNTRIES AND REGIONS

1. Bailey LB, Stover PJ, McNulty H, Fenech MF, Gregory JF, Mills JL, Pfeiffer CM, Fazili Z, Zhang M, Ueland PM, et al. Biomarkers of nutrition for development-Folate review. *J Nutr* 2015;145(7):1636S-80S.
2. World Health Organization. Serum and red blood cell folate concentrations for assessing folate status in populations. Vitamin and Mineral Nutrition Information System. Geneva: World Health Organization, 2015.
3. Pfeiffer CM, Lacher DA, Schleicher RL, Johnson CL, Yetley EA. Challenges and Lessons Learned in Generating and Interpreting NHANES Nutritional Biomarker Data. *Adv Nutr* 2017;8(2):290-307.
4. Pfeiffer CM, Fazili Z, Zhang M. Folate analytical methodology. Edition ed. In: Bailey LB, ed. *Folate in Health and Disease*. Boca Raton, FL: CRC Press, 2010:517-74.
5. Drammeh BS, Schleicher RL, Pfeiffer CM, Jain RB, Zhang M, Phuong HN. Effects of delayed sample processing and freezing on serum concentrations of selected nutritional indicators. *Clin Chem* 2008;54(11):1883-91.
6. O'Broin JD, Temperley IJ, Scott JM. Erythrocyte, plasma, and serum folate: specimen stability before microbiologic assay. *Clin Chem* 1980;26(3):522-4.
7. O'Broin SD, Kelleher BP, Davoren A, Gunter EW. Field-study screening of blood folate concentrations: Specimen stability and finger-stick sampling. *Am J Clin Nutr* 1997;66(6):1398-405.
8. Fazili Z, Sternberg MR, Pfeiffer CM. Assessing the influence of 5,10-methylenetetrahydrofolate reductase polymorphism on folate stability during long-term frozen storage, thawing, and repeated freeze/thawing of whole blood. *Clin Chim Acta* 2012;413(11-12):966-72.
9. Kelleher B, O'Broin S. Choice of materials for long-term quality control of blood folate assays. *Clin Chem* 1996;42(4):652-4.
10. O'Broin SD, Gunter EW. Screening of folate status with use of dried blood spots on filter paper. *Am J Clin Nutr* 1999;70(3):359-67.
11. Haynes BMH, Pfeiffer CM, Sternberg MR, Schleicher RL. Selected physiologic variables are weakly to moderately associated with 29 biomarkers of diet and nutrition, NHANES 2003-2006. *J Nutr* 2013;143(6):1001S-10S.
12. Gunter EW, Bowman BA, Caudill SP, Twite DB, Adams MJ, Sampson EJ. Results of an international round robin for serum and whole-blood folate. *Clin Chem* 1996;42(10):1689-94.
13. Pfeiffer CM, Gunter EW, Caudill SP. Comparison of serum and whole blood folate measurements in 12 laboratories: An international study. *Clin Chem* 2001;47:A62 (abstract 203).

14. Pfeiffer CM, Zhang M, Lacher DA, Molloy AM, Tamura T, Yetley EA, Picciano MF, Johnson CL. Comparison of serum and red blood cell folate microbiologic assays for national population surveys. *J Nutr* 2011;141(7):1402-9.
15. Bock JL, Endres DB, Elin RJ, Wang E, Rosenzweig B, Klee GG. Comparison of fresh frozen serum to traditional proficiency testing material in a College of American Pathologists survey for ferritin, folate, and vitamin B12. *Arch Pathol Lab Med* 2005;129(3):323-7.
16. Blackmore S, Pfeiffer CM, Lee A, Fazili Z, Hamilton MS. Isotope dilution-LC-MS/MS reference method assessment of serum folate assay accuracy and proficiency testing consensus mean. *Clin Chem* 2011;57(7):986-94.
17. Pfeiffer CM, Sternberg MR, Hamner HC, Crider KS, Lacher DA, Rogers LM, Bailey RL, Yetley EA. Applying inappropriate cutoffs leads to misinterpretation of folate status in the US population1-3. *Am J Clin Nutr* 2016;104(6):1607-15.
18. Institute of Medicine Standing Committee on the Scientific Evaluation of Dietary Reference Intakes and its Panel on Folate OBVaC. Dietary reference intakes for thamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin, and choline. Washington, DC: National Academie Press, 1998.
19. Selhub J, Jacques PF, Dallal G, Choumenkovitch S, Rogers G. The use of blood concentrations of vitamins and their respective functional indicators to define folate and vitamin B12 status. *Food Nutr Bull* 2008;29(2 SUPPL.).
20. World Health Organization. Guideline: Optimal Serum and Red Blood Cell Folate Concentrations in Women of Reproductive Age for Prevention of Neural Tube Defects. Geneva, Switzerland: World Health Organization, 2015.
21. Pfeiffer CM, Hughes JP, Durazo-Arvizu RA, Lacher DA, Sempos CT, Zhang M, Yetley EA, Johnson CL. Changes in measurement procedure from a radioassay to a microbiologic assay necessitate adjustment of serum and RBC folate concentrations in the U.S. population from the NHANES 1988-2010. *J Nutr* 2012;142(5):894-900.
22. Daly LE, Kirke PN, Molloy A, Weir DG, Scott JM. Folate Levels and Neural Tube Defects: Implications for Prevention. *JAMA* 1995;274(21):1698-702.
23. Tinker SC, Hamner HC, Qi YP, Crider KS. U.S. women of childbearing age who are at possible increased risk of a neural tube defect-affected pregnancy due to suboptimal red blood cell folate concentrations, National Health and Nutrition Examination Survey 2007 to 2012. *Birth Defects Research Part A - Clinical and Molecular Teratology* 2015;103(6):517-26.
24. Molloy AM, Scott JM. Microbiologic assay for serum, plasma, and red cell folate using cryopreserved, microtiter plate method. *Methods Enzymol* 1997;281:43-53.

A PUBLIC HEALTH APPROACH TO THE PREVENTION OF NTD: FORTIFICATION AND INNOVATIONS

1. Bailey L, et al. Biomarkers of Nutrition for Development – Folate Review. *J Nutr.* 2016.
2. Kyamuhangire W, Lubowa A, Kaaya A, Kikafunda J, et al. The importance of using food and nutrient intake data to identify appropriate vehicles and estimate potential benefits of food fortification in Uganda. *Food Nutr Bull.* 2013;34:131-142.
3. Czeizel AE, Dudas I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med.* 1992;327:1832-2835.
4. Centers for Disease Control and Prevention. Recommendations for the use of folic acid to reduce the number of cases of spina bifida and other neural tube defects. *Morbidity and Mortality Weekly Report (MMWR).* 1992;41(RR-14):1-7.
5. Berry RJ, Li Z, Erickson JD, et al. Prevention of neural-tube defects with folic acid in China. *N Engl J Med* 1999;341:1485-1890.
6. Honein MA, Paulozzi LJ, Mathews TJ, Erickson JD, Wong L-YC. Impact of folic acid fortification of the US food supply on the occurrence of neural tube defects. *JAMA.* 2001;285:2981-2986.
7. Ray JG, Wyatt PR, Thompson MD, et al. Vitamin B12 and the risk of neural tube defects in a folic-acid fortified population. *Epidemiology.* 2007;18:362-366.
8. Institute of Medicine. *Dietary Reference Intakes for thiamine, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin, and choline.* Washington (DC); 1988.
9. Global Panel of Agriculture and Food Systems for Nutrition. *Food systems and diets: Facing the challenges of the 21st century.* 2016.
10. Food Fortification Global Mapping Study. https://ec.europa.eu/europeaid/food-fortification-global-mapping-study-2016_en 2016.
11. Global Panel of Agriculture and Food Systems for Nutrition. “Food systems and diets: Facing the challenges of the 21st century”. <http://glopan.org/sites/default/files/ForesightReport.pdf2016>
12. Greiner T. Fortification of processed cereals should be mandatory. *Lancet.* 2007;9575: 1766-1768.
13. Fletcher R, Bell I, Lambert J. Public health aspects of food fortification: a question of balance. *Proc Nutr Soc.* 2004;63(4):605-614.
14. Bhutta Z, Das J, Rizwvi A, Gaffey M, Walker N, Horton S, et al. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? . *Lancet.* 2013;382(9890):452-477.
15. Healthbridge. *Fortification Rapid Assessment Tool and Guidelines.* 2003.
16. Neufeld LM, Dary O, Macdonald B, Harding KB. Measurement of food consumption to inform food fortification and other nutrition programs: Methods and Applications. *Food Nutr Bull.* 2012;23(Suppl. 3).

17. Neufeld L, Baker S, Garrett G, Haddad L. Coverage and utilization: Critical and neglected areas of program evaluation. *J Nutr*. 2016.
18. FAO, WHO. *Codex Alimentarius: general principles for the addition of essential nutrients to foods*. 2015.
19. Harvey P W, Dary O. Governments and academic institutions play vital roles in food fortification: iron as an example. *Public Health Nutr*. 2012;15(10):1791-1795.
20. Allen L, de Benoist B, Dary O, Hurrell R. *Guidelines on food fortification with micronutrients*. Geneva; 2006.
21. Honein MA, et al. Impact of folic acid fortification of the US food supply on the occurrence of neural tube defects. *JAMA*. 2001;285:2981–2986.
22. Country Profiles [Internet]. 2017 [cited 30 March 2017]. Available from: http://www.ffinetwork.org/country_profiles/data.php.
23. Food Fortification Initiative. *Plan for Fortification: Industry*. <http://www.ffinetwork.org/plan/industry.html> 2017.
24. Luthringer CL, et al. Regulatory monitoring of fortified foods: Identifying barriers and good practices. *Global Health Sci Pract*. 2015;3(3):446-461.
25. Aaron GJ, Friesen VM, Jungjohann S, Garrett GS, Neufeld LM, Myatt M. Coverage of large-scale food fortification of edible oil, wheat flour, and maize flour varies greatly by vehicle and country but is consistently lower among the most vulnerable: results from coverage surveys in 8 countries. *J Nutr*. 2017.
26. Cannon M. Preventing Neural Tube Defects (NTD) through Flour Fortification. *Global Summit on Food Fortification Plenary Presentation 2015*.
27. Youngblood ME, Williamson R, Bell KN, Johnson Q, Kancherla V, Oakley GP. 2012 Update on Global Prevention of Folic-Acid-Preventable Spina Bifida and Anencephaly. *Birth Defects Research (Part A) Clinic Molecular Teratol*. 2013;97:658-663.
28. CDC, Zaganjor, et al. unpublished data.
29. Crider , et al. *Br Med J*. 2014.
30. FFI , GAIN. *Feasibility and Potential Coverage of Fortified Rice in the Africa Rice Supply Chain*. 2016.
31. Li, al. e. Folic acid fortification through existing fortified foods: iodized salt and vitamin A-fortified sugar. *Food Nutr Bull*. 2011;32(1):35-41.
32. Mcgee E., et al. Double Fortification of Salt with Folic Acid and Iodine. *J Food Eng*. 2017;198:72-80.
33. WHO. *Salt reduction and iodine fortification strategies in public health*. 2013.
34. WHO. *Salt as a Vehicle for Fortification*. 2007.
35. Fortifying with Folic Acid [Internet]. [cited 24 April 2017]. Available from: <http://postgradasia.com/article/fortifying-with-folic-acid-1.aspx>.
36. Boy Erik. Personal communication. Head Nutrition, Harvest Plus; 30 March, 2017.

37. Global Panel of Agriculture and Food Systems for Nutrition. Food systems and diets: Facing the challenges of the 21st century. 2016
38. Storozhenko S, et al. Folate fortification of rice by metabolic engineering.
39. Blancquaert D, et al. Improving folate (vitamin B9) stability in biofortified rice through metabolic engineering. *Na Biotechnol.* 2015;33.
40. Kiekens F, et al. Folates from metabolically engineered rice: A long-term study in rats. *Mol Nutr Food Res.* 2015;59:490–500.
41. De Steur DS, De Steur H. Health impact assessment of folate biofortified rice in China. IDEAS Work Pap Ser from RePEc. 2010.
42. Lee H, Krimsky ST. The arrested development of Golden Rice: The Scientific and Social challenges of a transgenic biofortified crop. *Int J Soc Sci Stud.* 2016;4.
43. De Steur H, et al. Potential impact and cost-effectiveness of multi-biofortified rice in China. *N Biotechnol.* 2012;29:432–442

THE INVESTMENT CASE FOR FOLIC ACID FORTIFICATION

1. Castillo-Lancelotti C, et al. Impact of folic acid fortification of flour on neural tube defects: A systematic review. *Public Health Nutr.* 2013;16(5):901-911.
2. Sayed A-R, et al. Decline in the prevalence of neural tube defects following folic acid fortification and its cost benefit in South Africa. *Birth Defects Part A.* 2008;82:211-216.
3. Grosse S, et al. Retrospective assessment of cost savings from prevention: Folic acid fortification and Spina bifida in the US. *Am J Prev Med.* 2016;50(5S1):S74-S80.
4. Janneke J, et al. Economic evaluation of folic acid food fortification in The Netherlands. *Eur J Public Health.* 2008;18(3):270-274.
5. Access Economics. Cost benefit analysis of fortifying the food supply with folic acid. Canberra: Access Economics; 2006.
6. Health South Africa. Reflection on the South African maize meal and wheat flour fortification programme, 2004-2007. Pretoria: Health South Africa; 2008.
7. Berry R, et al. Fortification of flour with folic acid. *Food Nutr Bull.* 2010;31(1):S22-S35.
8. Prasad S. Folic acid prices and supply seen to ease up in 2016 and 2017. Accessed on February 8, 2017. Available from: <https://www.beroeinc.com/blog/folic-acid-prices-and-supply-seen-to-ease-up/>.
9. OECD. Development aid rises again in 2015, spending on refugees doubles. 2016 March 31, 2017. Available from: <http://www.oecd.org/dac/development-aid-rises-again-in-2015-spending-on-refugees-doubles.htm>
10. Statistics South Africa. Recorded live births 2014. Pretoria South Africa. 2015.
11. Rheingans, et al. Economic costs of rotavirus gastroenteritis and cost-effectiveness of vaccination in developing countries. *J Infect Dis.* 2009;200(Suppl 1):S16-S27.
12. Eisele T, et al. Estimates of child deaths prevented from malaria prevention scale-up in Africa 2001-2010. *Malaria J.* 2012;11(1):93-100.
13. Wang H, et al. Effectiveness of folic acid fortified flour for prevention of neural tube defects in a high risk region. *Nutrients.* 2016;8.
14. Llanos A, et al. Cost-effectiveness of a folic acid fortification program in Chile. *Health Affairs.* 2007;83(2-3):295-303.
15. Fiedler J, Lividini K. Managing the vitamin A program portfolio: A case study of Zambia, 2013-2042. *Food Nutr Bull.* 2014;35(1):105-125.
16. Horton S. The economics of food fortification. *J Nutr.* 2006;136(4):1068-1071.

17. Pulkki-Brannstrom A-M, et al. Cost and cost-effectiveness of long-lasting insecticide treated bed-nets: A model based analysis. *Cost effectiveness and resource allocation*. 2012;10(5).
18. Shekar M, et al. *An investment framework for nutrition*. Washington DC: World Bank; 2017.
19. Food Safety and Standards Authority of India (FSSAI). *Fortification of wheat flour: technology*. 2017 Accessed on April 15, 2017. Available from: <http://ffrc.fssai.gov.in/fortification/jsp/wheat.jsp>.
20. Hoddinott J, Gillespie S, Yosef S. *Public private partnerships and the reduction of undernutrition in developing countries*. Washington DC: International Food Policy Research Institute; 2015.
21. Aaron GJ, et al. Coverage of large-scale food fortification of edible oil, wheat flour, and maize flour varies greatly by vehicle and country but is consistently lower among the most vulnerable: Results from coverage surveys in 8 countries. *J Nutr*. 2017;147(5):S984-S994.
22. Luthringer C. Regulatory monitoring of fortified foods: Identifying barriers and good practices. *Global Health: Science and Practice*. 2015;3(3):446-461.
23. Hirvonen K, Hoddinott J. *Agricultural production and children's diets: Evidence from rural Ethiopia*. *Agricult Econ*. forthcoming.